arXiv:1107.1301v3 [math.RA] 24 Nov 2011

A CATALOG OF CAYLEY-DICKSON-LIKE PRODUCTS

JOHN W. BALES

Abstract. A catalog of all 32 Cayley-Dickson-like doubling products on ordered pairs $(a,b)\cdot(c,d)$ for which $(1,0)$ is the left and right identity and for which $x\cdot x^{*}=x^{*}\cdot x=\|x\|^{2}$ given the conjugate $(a,b)^{*}=(a^{*},-b)$. Only eight of these are true Cayley-Dickson doubling products, since 24 of them do not satisfy the quaternion properties. Each of the eight Cayley-Dickson products has a distinctive representation in the Fano Plane.

1. Product of finite sequences of real numbers

In this catalog, Cayley-Dickson-like products are considered to be products of finite sequences of real numbers. This may seem odd since Cayley-Dickson products are normally considered to be products defined on ordered pairs. In order to define a Cayley-Dickson-like product of any two finite sequences of real numbers, every finite sequence x will be identified with an infinite sequence of the form

$$x=x_0,x_1,x_2,\cdots,x_{n-1},0,0,0,\cdots$$

and the ordered pair of any two sequences x and y (whether finite or infinite) will be identified with the shuffling of the two sequences

$$(x,y)=x_0,y_0,x_1,y_1,x_2,y_2\cdots$$

If a is a real number, then a will be identified with the sequence $a,0,0,0,\cdots$.

The basis vectors $i_0,i_1,i_2,\cdots$ for this space are defined recursively.

$$\begin{aligned}i_0&=1\\ i_{2k}&=(i_k,0)\text{ for }k\geq 0\\ i_{2k+1}&=(0,i_k)\text{ for }k\geq 0\end{aligned}$$

Thus it follows that

Date: June 2011.

Key words and phrases. twisted group algebra, Cayley-Dickson algebra, quaternions, octonions, Fano plane.

$$\begin{aligned} i_0 &= 1,0,0,0,\cdots \\ i_1 &= 0,1,0,0,\cdots \\ i_2 &= 0,0,1,0,\cdots \\ &\vdots \end{aligned}$$

Furthermore, an involution * is defined on the basis vectors

$$i_k^* = \begin{cases} i_k & \text{if } k = 0 \\ -i_k & \text{if } k > 0 \end{cases}$$

and is extended to the entire space by the linearity property.

It then follows that for all infinite sequences x and y

$$(x,y)^* = (x^*, -y) \tag{1}$$

The norm $\|x\|$ of a finite sequence x is its Euclidean norm and the inner product $\langle x,y\rangle$ of two finite sequences x and y is their Euclidean inner product. Thus

$$\|(x,y)\|^2 = \|x\|^2 + \|y\|^2 = \left(\|x\|^2 + \|y\|^2, 0\right)$$

2. Cayley-Dickson-like Product

A Cayley-Dickson-like product on finite sequences (a,b) and (c,d) is a product satisfying the following four properties:

$$\begin{aligned} (1,0)(b,c) &= (b,c) && (2) \\ (a,b)(1,0) &= (a,b) && (3) \\ (a,b)(a^*,-b) &= \left(\|a\|^2 + \|b\|^2, 0\right) && (4) \\ (c^*,-d)(c,d) &= \left(\|c\|^2 + \|d\|^2, 0\right) && (5) \end{aligned}$$

It is 'Cayley-Dickson-like' rather than 'Cayley-Dickson' since we do not require the quaternion property.

The quaternion property is a property of the basis vectors. The statement that the quaternion property is satisfied means that if p,q and r are positive integers and if $i_p i_q = i_r$ then $i_q i_p = -i_r$ and $i_q i_r = i_p$.

The product of $(a,b)(c,d)$ can be defined 32 different ways to achieve the four 'Cayley-Dickson-like' properties, thus guaranteeing that for all finite sequences x,

$$1 \cdot x = x \cdot 1 = x \tag{6}$$

$$xx^* = x^*x = \|x\|^2 \tag{7}$$

Only eight of the 32 Cayley-Dickson-like products satisfy the quaternion properties. For all eight of these, $i_1 = i$. For four of them $i_2 = j$ and $i_3 = k$. For the other four, $i_2 = k$ and $i_3 = j$.

3. THE 32 WAYS

For each of the 32 Cayley-Dickson-like products of finite sequences, the recursive multiplication table for the basis vectors is given. The table is divided into distinct 2×2 blocks. These blocks occur either on the corner, left edge, top edge, diagonal or interior of the multiplication table.

An examination of the tables reveals that for all 32 variations it is true that $i_1 i_2 + i_2 i_1 = 0$. Using this as a basis step, it can be shown by induction that for all 32 variations

$$i_q i_p = -i_p i_q \text{ for } 0 \neq p \neq q \neq 0 \tag{8}$$

3.1. $P_0 : (a,b)(c,d) = (ca - b^*d, da^* + bc)$**.** For all r, s.

P_0	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r^*)$
i_{2r+1}	$(0, i_r i_s)$	$(-i_r^* i_s, 0)$

(1) For $r = s = 0$ (The corner)

P_0	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_0	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_0	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_0	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_0	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$-(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s)$	$(i_r i_s, 0)$

3.2. $P_1 : (a,b)(c,d) = (ca - db^*, da^* + bc)$**.** For all r, s.

P_1	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r^*)$
i_{2r+1}	$(0, i_r i_s)$	$(-i_s i_r^*, 0)$

(1) For $r = s = 0$ (The corner)

P_1	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_1	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_1	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_1	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_1	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$-(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s)$	$(i_s i_r, 0)$

3.3. $P_2 : (a,b)(c,d) = (ca - b^* d, a^* d + cb)$. For all r, s.

P_2	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r^* i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(-i_r^* i_s, 0)$

(1) For $r = s = 0$ (The corner)

P_2	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_2	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_2	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_2	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_2	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$-(0, i_r i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(i_r i_s, 0)$

3.4. **$P_3 : (a,b)(c,d) = (ca - db^*, a^*d + cb)$.** For all r, s.

P_3	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r^* i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(-i_s i_r^*, 0)$

(1) For $r = s = 0$ (The corner)

P_3	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_3	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_3	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_3	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_3	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$-(0, i_r i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(i_s i_r, 0)$

3.5. $P_4 : (a,b)(c,d) = (ac - b^*d, da^* + bc)$**.** For all r, s.

P_4	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r^*)$
i_{2r+1}	$(0, i_r i_s)$	$(-i_r^* i_s, 0)$

(1) For $r = s = 0$ (The corner)

P_4	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_4	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_4	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_4	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_4	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$-(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_r i_s, 0)$

3.6. $P_5 : (a,b)(c,d) = (ac - db^*, da^* + bc)$. For all r, s.

P_5	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r^*)$
i_{2r+1}	$(0, i_r i_s)$	$(-i_s i_r^*, 0)$

(1) For $r = s = 0$ (The corner)

P_5	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_5	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_5	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_5	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_5	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$-(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s)$	$(i_s i_r, 0)$

3.7. $P_6 : (a,b)(c,d) = (ac - b^*d, a^*d + cb)$**.** For all r, s.

P_6	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r^* i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(-i_r^* i_s, 0)$

(1) For $r = s = 0$ (The corner)

P_6	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_6	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_6	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_6	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_6	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$-(0, i_r i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(i_r i_s, 0)$

3.8. $P_7 : (a,b)(c,d) = (ac - db^*, a^*d + cb)$. For all r, s.

P_7	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r^* i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(-i_s i_r^*, 0)$

(1) For $r = s = 0$ (The corner)

P_7	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_7	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_7	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_7	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_7	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$-(0, i_r i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(i_s i_r, 0)$

3.9. $P_8 : (a,b)(c,d) = (ca - bd^*, da^* + bc)$**.** For all r, s.

P_8	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r^*)$
i_{2r+1}	$(0, i_r i_s)$	$(-i_r i_s^*, 0)$

(1) For $r = s = 0$ (The corner)

P_8	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_8	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_8	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_8	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_8	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$-(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s)$	$(i_r i_s, 0)$

3.10. $P_9 : (a,b)(c,d) = (ca - d^*b, da^* + bc)$**.** For all r, s.

P_9	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r^*)$
i_{2r+1}	$(0, i_r i_s)$	$(-i_s^* i_r, 0)$

(1) For $r = s = 0$ (The corner)

P_9	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_9	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_9	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_9	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_9	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_s, 0)$	$-(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s)$	$(i_s i_r, 0)$

3.11. $P_{10} : (a,b)(c,d) = (ca - bd^*, a^*d + cb)$**.** For all r, s.

P_{10}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r^* i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(-i_r i_s^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{10}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{10}	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{10}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{10}	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{10}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$-(0, i_r i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(i_r i_s, 0)$

3.12. $P_{11} : (a,b)(c,d) = (ca - d^*b, a^*d + cb)$**.** For all r, s.

P_{11}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r^* i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(-i_s^* i_r, 0)$

(1) For $r = s = 0$ (The corner)

P_{11}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{11}	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{11}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{11}	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{11}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$-(0, i_r i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(i_s i_r, 0)$

3.13. $P_{12} : (a,b)(c,d) = (ac - bd^*, da^* + bc)$**.** For all r, s.

P_{12}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r^*)$
i_{2r+1}	$(0, i_r i_s)$	$(-i_r i_s^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{12}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{12}	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{12}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{12}	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{12}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$-(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s)$	$(i_r i_s, 0)$

3.14. $P_{13} : (a,b)(c,d) = (ac - d^*b, da^* + bc)$**.** For all r, s.

P_{13}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r^*)$
i_{2r+1}	$(0, i_r i_s)$	$(-i_s^* i_r, 0)$

(1) For $r = s = 0$ (The corner)

P_{13}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{13}	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{13}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{13}	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{13}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$-(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s)$	$(i_s i_r, 0)$

3.15. $P_{14} : (a,b)(c,d) = (ac - bd^*, a^*d + cb)$**.** For all r, s.

P_{14}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r^* i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(-i_r i_s^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{14}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{14}	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{14}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{14}	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{14}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$-(0, i_r i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(i_r i_s, 0)$

3.16. $P_{15} : (a,b)(c,d) = (ac - d^*b, a^*d + cb)$**.** For all r, s.

P_{15}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r^* i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(-i_s^* i_r, 0)$

(1) For $r = s = 0$ (The corner)

P_{15}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{15}	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{15}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	i_{2s+1}	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{15}	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{15}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$-(0, i_r i_s)$
i_{2r+1}	$(0, i_s i_r)$	$(i_s i_r, 0)$

3.17. $P_{16} : (a,b)(c,d) = (ca - b^*d, ad + c^*b)$. For all r, s.

P_{16}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r i_s)$
i_{2r+1}	$(0, i_s^* i_r)$	$(-i_r^* i_s, 0)$

(1) For $r = s = 0$ (The corner)

P_{16}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{16}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{16}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_{16}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{16}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r i_s)$
i_{2r+1}	$-(0, i_s i_r)$	$(i_r i_s, 0)$

3.18. $P_{17} : (a, b)(c, d) = (ca - db^*, ad + c^*b)$**.** For all r, s.

P_{17}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r i_s)$
i_{2r+1}	$(0, i_s^* i_r)$	$(-i_s i_r^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{17}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{17}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{17}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_{17}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{17}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r i_s)$
i_{2r+1}	$-(0, i_s i_r)$	$(i_s i_r, 0)$

3.19. $P_{18} : (a, b)(c, d) = (ca - b^*d, da + bc^*)$**.** For all r, s.

P_{18}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s^*)d$	$(-i_r^* i_s, 0)$

(1) For $r = s = 0$ (The corner)

P_{18}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{18}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{18}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_{18}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{18}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_r i_s, 0)$

3.20. $P_{19} : (a,b)(c,d) = (ca - db^*, da + bc^*)$**.** For all r, s.

P_{19}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s^*)$	$(-i_s i_r^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{19}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{19}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{19}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_{19}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{19}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_s i_r, 0)$

3.21. $P_{20} : (a,b)(c,d) = (ac - b^*d, ad + c^*b)$**.** For all r, s.

P_{20}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r i_s)$
i_{2r+1}	$(0, i_s^* i_r)$	$(-i_r^* i_s, 0)$

(1) For $r = s = 0$ (The corner)

P_{20}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{20}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{20}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_{20}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{20}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r i_s)$
i_{2r+1}	$-(0, i_s i_r)$	$(i_r i_s, 0)$

3.22. $P_{21} : (a,b)(c,d) = (ac - db^*, ad + c^*b)$**.** For all r, s.

P_{21}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r i_s)$
i_{2r+1}	$(0, i_s^* i_r)$	$(-i_s i_r^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{21}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{21}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{21}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_{21}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{21}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r i_s)$
i_{2r+1}	$-(0, i_s i_r)$	$(i_s i_r, 0)$

3.23. $P_{22} : (a,b)(c,d) = (ac - b^*d, da + bc^*)$**.** For all r, s.

P_{22}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s^*)$	$(-i_r^* i_s, 0)$

(1) For $r = s = 0$ (The corner)

P_{22}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{22}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{22}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	$-i_{2s}$

(4) For $r = s > 0$ (The diagonal)

P_{22}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{22}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_r i_s, 0)$

3.24. $P_{23}:(a,b)(c,d)=(ac-db^*,da+bc^*)$**.** For all r,s.

P_{23}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s^*)$	$(-i_s i_r^*, 0)$

(1) For $r=s=0$ (The corner)

P_{23}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r>s=0$ (The left edge)

P_{23}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s>r=0$ (The top edge)

P_{23}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	$-i_{2s}$

(4) For $r=s>0$ (The diagonal)

P_{23}	i_{2r}	i_{2r+1}
i_{2r}	-1	i_1
i_{2r+1}	$-i_1$	-1

(5) For $0\neq r\neq s\neq 0$ (The interior)

P_{23}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_s i_r, 0)$

3.25. $P_{24}:(a,b)(c,d)=(ca-bd^*,ad+c^*b)$**.** For all r,s.

P_{24}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r i_s)$
i_{2r+1}	$(0, i_s^* i_r)$	$(-i_r i_s^*, 0)$

(1) For $r=s=0$ (The corner)

P_{24}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r>s=0$ (The left edge)

P_{24}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s>r=0$ (The top edge)

P_{24}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	i_{2s}

(4) For $r=s>0$ (The diagonal)

P_{24}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0\neq r\neq s\neq 0$ (The interior)

P_{24}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r i_s)$
i_{2r+1}	$-(0, i_s i_r)$	$(i_r i_s, 0)$

3.26. $P_{25} : (a,b)(c,d) = (ca - d^*b, ad + c^*b)$**.** For all r, s.

P_{25}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r i_s)$
i_{2r+1}	$(0, i_s^* i_r)$	$(-i_s^* i_r, 0)$

(1) For $r = s = 0$ (The corner)

P_{25}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{25}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{25}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{25}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{25}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_r i_s)$
i_{2r+1}	$-(0, i_s i_r)$	$(i_s i_r, 0)$

3.27. $P_{26} : (a,b)(c,d) = (ca - bd^*, da + bc^*)$**.** For all r, s.

P_{26}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s^*)$	$(-i_r i_s^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{26}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{26}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{26}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{26}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{26}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_r i_s, 0)$

3.28. $P_{27} : (a,b)(c,d) = (ca - d^*b, da + bc^*)$. For all r, s.

P_{27}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s^*)$	$(-i_s^* i_r, 0)$

(1) For $r = s = 0$ (The corner)

P_{27}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{27}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{27}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{27}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{27}	i_{2s}	i_{2s+1}
i_{2r}	$(i_s i_r, 0)$	$(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_s i_r, 0)$

3.29. $P_{28} : (a, b)(c, d) = (ac - bd^*, ad + c^*b)$. For all r, s.

P_{28}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r i_s)$
i_{2r+1}	$(0, i_s^* i_r)$	$(-i_r i_s^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{28}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{28}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{28}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{28}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{28}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r i_s)$
i_{2r+1}	$-(0, i_s i_r)$	$(i_r i_s, 0)$

3.30. $P_{29} : (a,b)(c,d) = (ac - d^*b, ad + c^*b)$**.** For all r, s.

P_{29}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r i_s)$
i_{2r+1}	$(0, i_s^* i_r)$	$(-i_s^* i_r, 0)$

(1) For $r = s = 0$ (The corner)

P_{29}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{29}	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{29}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{29}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{29}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_r i_s)$
i_{2r+1}	$-(0, i_s i_r)$	$(i_s i_r, 0)$

3.31. $P_{30} : (a, b)(c, d) = (ac - bd^*, da + bc^*)$. For all r, s.

P_{30}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s^*)$	$(-i_r i_s^*, 0)$

(1) For $r = s = 0$ (The corner)

P_{30}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{30}	i_0	i_1
i_{2r}	i_{2r}	i_{2r+1}
i_{2r+1}	i_{2r+1}	$-i_{2r}$

(3) For $s > r = 0$ (The top edge)

P_{30}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_90	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{30}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_r i_s, 0)$

3.32. $P_{31} : (a,b)(c,d) = (ac - d^*b, da + bc^*)$**.** For all r, s.

P_{31}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r)$
i_{2r+1}	$(0, i_r i_s^*)$	$(-i_s^* i_r, 0)$

(1) For $r = s = 0$ (The corner)

P_{31}	i_0	i_1
i_0	1	i_1
i_1	i_1	-1

(2) For $r > s = 0$ (The left edge)

P_{31}	i_0	i_1
i_{2r}	i_{2r}	$-i_{2r+1}$
i_{2r+1}	i_{2r+1}	i_{2r}

(3) For $s > r = 0$ (The top edge)

P_{31}	i_{2s}	i_{2s+1}
i_0	i_{2s}	i_{2s+1}
i_1	$-i_{2s+1}$	i_{2s}

(4) For $r = s > 0$ (The diagonal)

P_{31}	i_{2r}	i_{2r+1}
i_{2r}	-1	$-i_1$
i_{2r+1}	i_1	-1

(5) For $0 \neq r \neq s \neq 0$ (The interior)

P_{31}	i_{2s}	i_{2s+1}
i_{2r}	$(i_r i_s, 0)$	$(0, i_s i_r)$
i_{2r+1}	$-(0, i_r i_s)$	$(i_s i_r, 0)$

4. CAYLEY-DICKSON-LIKE PRODUCTS AS TWISTED GROUP PRODUCTS

The basis elements $i_0, i_1, i_2, \cdots$ are indexed by $W = \{0, 1, 2, \cdots\}$ which is a group under the bit-wise 'exclusive or' of their binary representations and for each of the 32 Cayley-Dickson-like products there is a function ω from $W \times W$ to $\{-1, 1\}$ such that for $p, q \in W$

$$i_p i_q = \omega(p, q) i_{pq} \tag{9}$$

where pq is the group product of p and q. The function ω is called a 'twist' on the group W and turns the set of all finite sequences into a twisted group algebra.

Since $1 = (1, 0) = i_0$ is the identity, it follows that $\omega(p, 0) = \omega(0, q) = 1$ for all Cayley-Dickson-like products, and since for $p > 1$, $i_p i_p^* = \|i_p\|^2 = 1$ and since $i_p i_p^* = -i_p i_p = -\omega(p, p) i_0 = 1$ it follows that for $p > 1$

$$\omega(p, p) = -1 \tag{10}$$

and that for all p

$$i_p^* = \omega(p, p) i_p \tag{11}$$

From equation 8 on page 3 it follows that

$$\omega(q, p) + \omega(p, q) = 0 \text{ for } 0 \neq p \neq q \neq 0 \tag{12}$$

4.1. The product of $i_{2r} i_{2s}$.

Since $i_{2r} = (i_r, 0)$ then $i_{2r} i_{2s} = (i_r, 0)\,(i_s, 0)$ this could be called the case of $b = d = 0$. For each of the 32 distinct products, either $i_{2r} i_{2s} = (i_s i_r, 0)$ or $i_{2r} i_{2s} = (i_r i_s, 0)$. We shall consider the effect each of these alternatives upon the twist ω.

4.1.1. $i_{2r} i_{2s} = (i_s i_r, 0)$.

Since $i_{2r} i_{2s} = \omega(2r, 2s) i_{2rs}$ and since $i_{2r} i_{2s} = \omega(s, r) i_{2rs}$ we may conclude that Whenever $i_{2r} i_{2s} = (i_s i_r, 0)$,

$$\omega(2r, 2s) = \omega(s, r) \tag{13}$$

4.1.2. $i_{2r} i_{2s} = (i_r i_s, 0)$.

Since $i_{2r} i_{2s} = \omega(2r, 2s) i_{2rs}$ and since $i_{2r} i_{2s} = \omega(r, s) i_{2rs}$ we may conclude that Whenever $i_{2r} i_{2s} = (i_r i_s, 0)$,

$$\omega(2r, 2s) = \omega(r, s) \tag{14}$$

4.2. **The products of $i_{2r}i_{2s+1}$ and $i_{2r+1}i_{2s}$.** Since $i_{2r}i_{2s+1} = (i_r, 0)\,(0, i_s)$ this could be called the case of $b = c = 0$. And since $i_{2r+1}i_{2s} = (0, i_r)\,(i_s, 0)$ that could be called the case of $a = d = 0$. An inspection of the 32 alternate products shows that there are only four distinct possibilities for the products of $i_{2r}i_{2s+1}$ and $i_{2r+1}i_{2s}$.

4.2.1. *$i_{2r}i_{2s+1} = (0, i_s i_r)$ and $i_{2r+1}i_{2s} = (0, i_r i_s^*)$.* These two conditions imply that whenever $i_{2r}i_{2s+1} = (0, i_s i_r)$,

$$\omega(2r, 2s+1) = \omega(s, r) \tag{15}$$

and that whenever $i_{2r+1}i_{2s} = (i_r i_s^*, 0)$,

$$\omega(2r+1, 2s) = \omega(s, s)\omega(r, s) = \begin{cases} -\omega(r, s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases} \tag{16}$$

4.2.2. *$i_{2r}i_{2s+1} = (0, i_r i_s)$ and $i_{2r+1}i_{2s} = (0, i_s^* i_r)$.* These two conditions imply that whenever $i_{2r}i_{2s+1} = (0, i_r i_s)$,

$$\omega(2r, 2s+1) = \omega(r, s) \tag{17}$$

and that whenever $i_{2r+1}i_{2s} = (i_s^* i_r, 0)$,

$$\omega(2r+1, 2s) = \omega(s, s)\omega(s, r) = \begin{cases} -\omega(s, r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases} \tag{18}$$

4.2.3. *$i_{2r}i_{2s+1} = (0, i_s i_r^*)$ and $i_{2r+1}i_{2s} = (0, i_r i_s)$.* These two conditions imply that whenever $i_{2r}i_{2s+1} = (0, i_s i_r^*)$

$$\omega(2r, 2s+1) = \omega(r, r)\omega(s, r) = \begin{cases} -\omega(s, r) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases} \tag{19}$$

and that whenever $i_{2r+1}i_{2s} = (i_r i_s, 0)$,

$$\omega(2r+1, 2s) = \omega(r, s) \tag{20}$$

4.2.4. *$i_{2r}i_{2s+1} = (0, i_r^* i_s)$ and $i_{2r+1}i_{2s} = (0, i_s i_r)$.* These two conditions imply that whenever $i_{2r}i_{2s+1} = (0, i_r^* i_s)$

$$\omega(2r, 2s+1) = \omega(s, s)\omega(r, s) = \begin{cases} -\omega(r, s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases} \tag{21}$$

and that whenever $i_{2r+1}i_{2s} = (i_s i_r, 0)$,

$$\omega(2r+1,2s)=\omega(r,s) \tag{22}$$

4.3. **The product of $i_{2r+1}i_{2s+1}$.** Since $i_{2r+1}=(0,i_r)$ the product $i_{2r+1}i_{2s+1}$ could be called the case of $a=c=0$.

4.3.1. $i_{2r+1}i_{2s+1}=-\left(i_s^* i_r,0\right)$. This implies that whenever $i_{2r+1}i_{2s+1}=-\left(i_s^* i_r,0\right)$

$$\omega(2r+1,2s+1)=-\omega(s,s)\omega(s,r)=\begin{cases}\omega(s,r) \text{ if } s>0\\ -1 \text{ otherwise}\end{cases} \tag{23}$$

4.3.2. $i_{2r+1}i_{2s+1}=-\left(i_r i_s^*,0\right)$. This implies that whenever $i_{2r+1}i_{2s+1}=-\left(i_r i_s^*,0\right))$

$$\omega(2r+1,2s+1)=-\omega(s,s)\omega(r,s)=\begin{cases}\omega(r,s) \text{ if } s>0\\ -1 \text{ otherwise}\end{cases} \tag{24}$$

4.3.3. $i_{2r+1}i_{2s+1}=-\left(i_s i_r^*,0\right)$. This implies that whenever $i_{2r+1}i_{2s+1}=-\left(i_s i_r^*,0\right)$

$$\omega(2r+1,2s+1)=-\omega(r,r)\omega(s,r)=\begin{cases}\omega(s,r) \text{ if } r>0\\ -1 \text{ otherwise}\end{cases} \tag{25}$$

4.3.4. $i_{2r+1}i_{2s+1}=-\left(i_r^* i_s,0\right)$. This implies that whenever $i_{2r+1}i_{2s+1}=-\left(i_r^* i_s,0\right)$

$$\omega(2r+1,2s+1)=-\omega(r,r)\omega(r,s)=\begin{cases}\omega(r,s) \text{ if } r>0\\ -1 \text{ otherwise}\end{cases} \tag{26}$$

5. THE 32 VARIATIONS ON ω

For each of the 32 Cayley-Dickson-like products of finite sequences, the recursive definitions for ω are given.

(0) $\boxed{P_0 : (a,b)(c,d) = (ca - b^*d, da^* + bc)}$

ω_0	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\begin{cases} -\omega(s,r) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(r,s)$	$\begin{cases} \omega(r,s) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

(1) $\boxed{P_1 : (a,b)(c,d) = (ca - db^*, da^* + bc)}$

ω_1	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\begin{cases} -\omega(s,r) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(r,s)$	$\begin{cases} \omega(s,r) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

(2) $\boxed{P_2 : (a,b)(c,d) = (ca - b^*d, a^*d + cb)}$

ω_2	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\begin{cases} -\omega(r,s) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(s,r)$	$\begin{cases} \omega(r,s) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

(3) $\boxed{P_3 : (a,b)(c,d) = (ca - db^*, a^*d + cb)}$

ω_3	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\begin{cases} -\omega(r,s) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(s,r)$	$\begin{cases} \omega(s,r) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_4 : (a,b)(c,d) = (ac - b^*d, da^* + bc)} \tag{4}$$

ω_4	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\begin{cases} -\omega(s,r) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(r,s)$	$\begin{cases} \omega(r,s) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_5 : (a,b)(c,d) = (ac - db^*, da^* + bc)} \tag{5}$$

ω_5	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\begin{cases} -\omega(s,r) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(r,s)$	$\begin{cases} \omega(s,r) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_6 : (a,b)(c,d) = (ac - b^*d, a^*d + cb)} \tag{6}$$

ω_6	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\begin{cases} -\omega(r,s) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(s,r)$	$\begin{cases} \omega(r,s) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_7 : (a,b)(c,d) = (ac - db^*, a^*d + cb)} \tag{7}$$

ω_7	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\begin{cases} -\omega(r,s) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(s,r)$	$\begin{cases} \omega(s,r) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_8 : (a,b)(c,d) = (ca - bd^*, da^* + bc)} \tag{8}$$

ω_8	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\begin{cases} -\omega(s,r) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(r,s)$	$\begin{cases} \omega(r,s) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_9 : (a,b)(c,d) = (ca - d^*b, da^* + bc)} \tag{9}$$

ω_9	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\begin{cases} -\omega(s,r) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(r,s)$	$\begin{cases} \omega(s,r) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{10} : (a,b)(c,d) = (ca - bd^*, a^*d + cb)} \tag{10}$$

ω_{10}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\begin{cases} -\omega(r,s) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(s,r)$	$\begin{cases} \omega(r,s) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{11} : (a,b)(c,d) = (ca - d^*b, a^*d + cb)} \tag{11}$$

ω_{11}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\begin{cases} -\omega(r,s) \text{ if } r>0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(s,r)$	$\begin{cases} \omega(s,r) \text{ if } s>0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{12} : (a,b)(c,d) = (ac - bd^*, da^* + bc)} \tag{12}$$

ω_{12}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\begin{cases} -\omega(s,r) \text{ if } r>0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(r,s)$	$\begin{cases} \omega(r,s) \text{ if } s>0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{13} : (a,b)(c,d) = (ac - d^*b, da^* + bc)} \tag{13}$$

ω_{13}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\begin{cases} -\omega(s,r) \text{ if } r>0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(r,s)$	$\begin{cases} \omega(s,r) \text{ if } s>0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{14} : (a,b)(c,d) = (ac - bd^*, a^*d + cb)} \tag{14}$$

ω_{14}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\begin{cases} -\omega(r,s) \text{ if } r>0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(s,r)$	$\begin{cases} \omega(r,s) \text{ if } s>0 \\ -1 \text{ otherwise} \end{cases}$

(15) $\boxed{P_{15} : (a,b)(c,d) = (ac - d^*b, a^*d + cb)}$

ω_{15}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\begin{cases} -\omega(r,s) \text{ if } r > 0 \\ 1 \text{ otherwise} \end{cases}$
$2r+1$	$\omega(s,r)$	$\begin{cases} \omega(s,r) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

(16) $\boxed{P_{16} : (a,b)(c,d) = (ca - b^*d, ad + c^*b)}$

ω_{16}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\omega(r,s)$
$2r+1$	$\begin{cases} -\omega(s,r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(r,s) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

(17) $\boxed{P_{17} : (a,b)(c,d) = (ca - db^*, ad + c^*b)}$

ω_{17}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\omega(r,s)$
$2r+1$	$\begin{cases} -\omega(s,r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(s,r) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

(18) $\boxed{P_{18} : (a,b)(c,d) = (ca - b^*d, da + bc^*)}$

ω_{18}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\omega(s,r)$
$2r+1$	$\begin{cases} -\omega(r,s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(r,s) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

(19) $\boxed{P_{19} : (a,b)(c,d) = (ca - db^*, da + bc^*)}$

ω_{19}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\omega(s,r)$
$2r+1$	$\begin{cases} -\omega(r,s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(s,r) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{20} : (a,b)(c,d) = (ac - b^*d, ad + c^*b)} \tag{20}$$

ω_{20}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\omega(r,s)$
$2r+1$	$\begin{cases} -\omega(s,r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(r,s) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{21} : (a,b)(c,d) = (ac - db^*, ad + c^*b)} \tag{21}$$

ω_{21}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\omega(r,s)$
$2r+1$	$\begin{cases} -\omega(s,r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(s,r) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{22} : (a,b)(c,d) = (ac - b^*d, da + bc^*)} \tag{22}$$

ω_{22}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\omega(s,r)$
$2r+1$	$\begin{cases} -\omega(r,s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(r,s) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{23} : (a,b)(c,d) = (ac - db^*, da + bc^*)} \tag{23}$$

ω_{23}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\omega(s,r)$
$2r+1$	$\begin{cases} -\omega(r,s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(s,r) \text{ if } r > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{24} : (a,b)(c,d) = (ca - bd^*, ad + c^*b)} \tag{24}$$

ω_{24}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\omega(r,s)$
$2r+1$	$\begin{cases} -\omega(s,r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(r,s) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{25} : (a,b)(c,d) = (ca - d^*b, ad + c^*b)} \tag{25}$$

ω_{25}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\omega(r,s)$
$2r+1$	$\begin{cases} -\omega(s,r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(s,r) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{26} : (a,b)(c,d) = (ca - bd^*, da + bc^*)} \tag{26}$$

ω_{26}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\omega(s,r)$
$2r+1$	$\begin{cases} -\omega(r,s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(r,s) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{27} : (a,b)(c,d) = (ca - d^*b, da + bc^*)} \tag{27}$$

ω_{27}	$2s$	$2s+1$
$2r$	$\omega(s,r)$	$\omega(s,r)$
$2r+1$	$\begin{cases} -\omega(r,s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(s,r) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{28} : (a,b)(c,d) = (ac - bd^*, ad + c^*b)} \tag{28}$$

ω_{28}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\omega(r,s)$
$2r+1$	$\begin{cases} -\omega(s,r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(r,s) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{29} : (a,b)(c,d) = (ac - d^*b, ad + c^*b)} \tag{29}$$

ω_{29}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\omega(r,s)$
$2r+1$	$\begin{cases} -\omega(s,r) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(s,r) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{30} : (a,b)(c,d) = (ac - bd^*, da + bc^*)} \tag{30}$$

ω_{30}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\omega(s,r)$
$2r+1$	$\begin{cases} -\omega(r,s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(r,s) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

$$\boxed{P_{31} : (a,b)(c,d) = (ac - d^*b, da + bc^*)} \tag{31}$$

ω_{31}	$2s$	$2s+1$
$2r$	$\omega(r,s)$	$\omega(s,r)$
$2r+1$	$\begin{cases} -\omega(r,s) \text{ if } s > 0 \\ 1 \text{ otherwise} \end{cases}$	$\begin{cases} \omega(s,r) \text{ if } s > 0 \\ -1 \text{ otherwise} \end{cases}$

6. THE TWIST BLOCKS

For each of the 32 twists ω_k on the group $\{0, 1, 2, 3, \cdots\}$ there are five fundamental 2×2 matrices $E_{krs} = \begin{bmatrix} \omega_k(2r, 2s) & \omega_k(2r, 2s+1) \\ \omega_k(2r+1, 2s) & \omega_k(2r+1, 2s+1) \end{bmatrix}$ which make up the twist table.

- For $r = s = 0$, $E_{k00} = \omega_k(0,0)C$, where C is the block in the upper left corner of the twist table.
- For $r > s = 0$, $E_{kr0} = \omega_k(r,0)L$, where L is the block occupying every position in the left edge of the twist table below the corner block.
- For $s > r = 0$, $E_{k0s} = \omega_k(0,s)T$, where T is the block occupying every position in the top edge of the twist table to the right of the corner block.
- For $r = s \neq 0$, $E_{krr} = \omega_k(r,r)D$, where D is the block whose negative occupies every position on the diagonal of the twist table with the exception of the corner block.
- For $0 \neq r \neq s \neq 0$, $E_{krs} = \omega_k(r,s)N$, where N is the interior block of the twist table. Either N or its negative (depending upon the value of $\omega(r,s)$) occupies the twist table in block position (r,s) (counting the first block row as block row zero and the first block column as block column 0) when $0 \neq r \neq s \neq 0$.

The values of the five fundamental blocks $\mathbf{C}, \mathbf{L}, \mathbf{T}, \mathbf{D}$ and $\mathbf{N}$ independent of the value of $\omega(r,s)$ are given for each of the 32 twists in Table 1 and Table 2.

Twist	**C**	**L**	**T**	**D**	**N**
	$r = s = 0$	$r > s = 0$	$s > r = 0$	$r = s \neq 0$	$0 \neq r \neq s \neq 0$
ω_0	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix}$
ω_1	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}$
ω_2	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix}$
ω_3	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$
ω_4	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$
ω_5	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$
ω_6	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$
ω_7	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix}$
ω_8	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix}$
ω_9	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}$
ω_{10}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix}$
ω_{11}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$
ω_{12}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$
ω_{13}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$
ω_{14}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$
ω_{15}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix}$

TABLE 1. Twist blocks for ω_0 through ω_{15}

Twist	**C**	**L**	**T**	**D**	**N**
	$r = s = 0$	$r > s = 0$	$s > r = 0$	$r = s \neq 0$	$0 \neq r \neq s \neq 0$
ω_{16}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix}$
ω_{17}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}$
ω_{18}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix}$
ω_{19}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$
ω_{20}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$
ω_{21}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$
ω_{22}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$
ω_{23}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix}$
ω_{24}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 1 \\ 1 & 1 \end{bmatrix}$
ω_{25}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & 1 \\ 1 & -1 \end{bmatrix}$
ω_{26}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ -1 & 1 \end{bmatrix}$
ω_{27}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} -1 & -1 \\ -1 & -1 \end{bmatrix}$
ω_{28}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$
ω_{29}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$
ω_{30}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ -1 & 1 \end{bmatrix}$
ω_{31}	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ -1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ -1 & -1 \end{bmatrix}$

TABLE 2. Twist blocks for ω_{16} through ω_{31}

7. The Twist Tree

Looking at tables 1 and 2 it is seen that the five basic twist blocks are:

$$\mathbf{C} = \begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix} \tag{27}$$

$$\mathbf{L} = \begin{bmatrix} 1 & \alpha' \\ 1 & \alpha \end{bmatrix} \tag{28}$$

$$\mathbf{T} = \begin{bmatrix} 1 & 1 \\ \alpha & \alpha' \end{bmatrix} \tag{29}$$

$$\mathbf{D} = \begin{bmatrix} 1 & \alpha' \\ \alpha & 1 \end{bmatrix} \tag{30}$$

$$\mathbf{N} = \begin{bmatrix} \beta & \gamma \\ \gamma & \gamma' \end{bmatrix} \tag{31}$$

The five twist constants $\alpha', \alpha, \beta, \gamma, \gamma'$ are always equal to either 1 or -1.

Square matrices having 2^N rows and columns can be partitioned into 2×2 partitioned block matrices as follows:

$$M = \left[\begin{array}{c:c} M_{0,0} & M_{0,1} \\ \hdashline M_{1,0} & M_{1,1} \end{array}\right] \tag{32}$$

Each submatrix, provided it is not a 1×1 matrix, can be further subdivided. For example

$$M_{1,0} = \left[\begin{array}{c:c} M_{10,00} & M_{10,01} \\ \hdashline M_{11,00} & M_{11,01} \end{array}\right] \tag{33}$$

Then if the rows and columns of the matrices are numbered beginning with 0 rather than 1 and when, ultimately $M_{p,q}$ is a 1×1 matrix, it will be the case that

$$M_{i,j} = m_{ij} \tag{34}$$

Equation 32 can be written in 'tree' form as follows:



The subscripts can be viewed as navigation instructions: A *zero* is an instruction to move down a left branch and a *one* is an instruction to move down a right branch of the tree. Each of the block matrices $\mathbf{C}, \mathbf{L}, \mathbf{T}, \mathbf{D}$ and $\mathbf{N}$ can be written in tree form.

The twist tree for σ_7 was derived in [4]. Table 3 shows the general twist tree for all 32 variations on the Cayley-Dickson doubling formula.



Table 3. Generalized ‘Cayley-Dickson-Like’ Twist Tree

The values of $\alpha, \alpha', \beta, \gamma$ and γ' for the 32 different doublings are given in table 4 on the next page.

8. The quaternion properties

The two quaternion properties, stated in terms of the twist ω are as follows:

If $0 \neq p \neq q \neq 0$ then

(1) $\omega(p, q) + \omega(q, p) = 0$

(2) $\omega(p, q) = \omega(q, pq) = \omega(pq, p)$

$\omega(q, p) + \omega(p, q) = 0$ was previously shown to be true for all 32 products (see equation 12 on page 36).

	α'	α	β	γ	γ'
$P_0 : (a,b)(c,d) = (ca - b^*d, da^* + bc)$	$-$	$+$	$-$	$+$	$+$
$P_1 : (a,b)(c,d) = (ca - db^*, da^* + bc)$	$-$	$+$	$-$	$+$	$-$
$P_2 : (a,b)(c,d) = (ca - b^*d, a^*d + cb)$	$-$	$+$	$-$	$-$	$+$
$P_3 : (a,b)(c,d) = (ca - db^*, a^*d + cb)$	$-$	$+$	$-$	$-$	$-$
$P_4 : (a,b)(c,d) = (ac - b^*d, da^* + bc)$	$-$	$+$	$+$	$+$	$+$
$P_5 : (a,b)(c,d) = (ac - db^*, da^* + bc)$	$-$	$+$	$+$	$+$	$-$
$P_6 : (a,b)(c,d) = (ac - b^*d, a^*d + cb)$	$-$	$+$	$+$	$-$	$+$
$P_7 : (a,b)(c,d) = (ac - db^*, a^*d + cb)$	$-$	$+$	$+$	$-$	$-$
$P_8 : (a,b)(c,d) = (ca - bd^*, da^* + bc)$	$-$	$-$	$-$	$+$	$+$
$P_9 : (a,b)(c,d) = (ca - d^*b, da^* + bc)$	$-$	$-$	$-$	$+$	$-$
$P_{10} : (a,b)(c,d) = (ca - bd^*, a^*d + cb)$	$-$	$-$	$-$	$-$	$+$
$P_{11} : (a,b)(c,d) = (ca - d^*b, a^*d + cb)$	$-$	$-$	$-$	$-$	$-$
$P_{12} : (a,b)(c,d) = (ac - bd^*, da^* + bc)$	$-$	$-$	$+$	$+$	$+$
$P_{13} : (a,b)(c,d) = (ac - d^*b, da^* + bc)$	$-$	$-$	$+$	$+$	$-$
$P_{14} : (a,b)(c,d) = (ac - bd^*, a^*d + cb)$	$-$	$-$	$+$	$-$	$+$
$P_{15} : (a,b)(c,d) = (ac - d^*b, a^*d + cb)$	$-$	$-$	$+$	$-$	$-$
$P_{16} : (a,b)(c,d) = (ca - b^*d, ad + c^*b)$	$+$	$+$	$-$	$+$	$+$
$P_{17} : (a,b)(c,d) = (ca - db^*, ad + c^*b)$	$+$	$+$	$-$	$+$	$-$
$P_{18} : (a,b)(c,d) = (ca - b^*d, da + bc^*)$	$+$	$+$	$-$	$-$	$+$
$P_{19} : (a,b)(c,d) = (ca - db^*, da + bc^*)$	$+$	$+$	$-$	$-$	$-$
$P_{20} : (a,b)(c,d) = (ac - b^*d, ad + c^*b)$	$+$	$+$	$+$	$+$	$+$
$P_{21} : (a,b)(c,d) = (ac - db^*, ad + c^*b)$	$+$	$+$	$+$	$+$	$-$
$P_{22} : (a,b)(c,d) = (ac - b^*d, da + bc^*)$	$+$	$+$	$+$	$-$	$+$
$P_{23} : (a,b)(c,d) = (ac - db^*, da + bc^*)$	$+$	$+$	$+$	$-$	$-$
$P_{24} : (a,b)(c,d) = (ca - bd^*, ad + c^*b)$	$+$	$-$	$-$	$+$	$+$
$P_{25} : (a,b)(c,d) = (ca - d^*b, ad + c^*b)$	$+$	$-$	$-$	$+$	$-$
$P_{26} : (a,b)(c,d) = (ca - bd^*, da + bc^*)$	$+$	$-$	$-$	$-$	$+$
$P_{27} : (a,b)(c,d) = (ca - d^*b, da + bc^*)$	$+$	$-$	$-$	$-$	$-$
$P_{28} : (a,b)(c,d) = (ac - bd^*, ad + c^*b)$	$+$	$-$	$+$	$+$	$+$
$P_{29} : (a,b)(c,d) = (ac - d^*b, ad + c^*b)$	$+$	$-$	$+$	$+$	$-$
$P_{30} : (a,b)(c,d) = (ac - bd^*, da + bc^*)$	$+$	$-$	$+$	$-$	$+$
$P_{31} : (a,b)(c,d) = (ac - d^*b, da + bc^*)$	$+$	$-$	$+$	$-$	$-$

TABLE 4. Constants for the 32 doubling products

It will be shown that $\omega(p,q) = \omega(q,pq) = \omega(pq,p)$ for only eight of the 32 Cayley-Dickson-like doubling products. These eight are the only actual Cayley-Dickson products.

8.1. **The basis step of the induction.** Begin by showing that the property holds when either p or q equals 1 or when p and q differ by 1. This forms the basis step of an induction.

For $p = 2r \neq 0$ and $q = 1$ we have $pq = 2r + 1$, so we must compare the values of $\omega(2r, 1), \omega(1, 2r + 1)$ and $\omega(2r + 1, 2r)$.

$\omega(2r, 1) = \omega(2r, 2(0) + 1)$ so we find the value in the $(2r, 2s + 1)$ portions of the charts with $r > 0, s = 0$. We see that the value is -1 for ω_0 through ω_{15} and 1 for ω_{16} through ω_{31}.

$\omega(1, 2r + 1) = \omega(2(0) + 1, 2s + 1)$ We find these values in the $(2r + 1, 2s + 1)$ portion of the tables with $r = 0$ and $s > 0$. We see that the value is -1 for ω_0 through ω_7 and ω_{16} through ω_{23} and 1 for ω_8 through ω_{15} and ω_{24} through ω_{31}.

We conclude that the twists ω_8 through ω_{23} fail to have the second quaternion property.

$\omega(2r, 2r + 1)$ has a value of -1 for ω_0 through ω_{15} and 1 for ω_{16} through ω_{31}.

Thus $\omega(2r, 1) = \omega(1, 2r + 1) = \omega(2r + 1, 2r)$ for ω_0 through ω_7 and for ω_{24} through ω_{31} alone.

Since $\omega(2r, 1) = -\omega(1, 2r), \omega(1, 2r + 1) = -\omega(2r + 1, 1)$ and $\omega(2r + 1, 2r) = -\omega(2r, 2r+1)$ we have the result that $\omega(2r+1, 1) = \omega(1, 2r) = \omega(2r, 2r + 1)$.

So for ω_0 through ω_7 and for ω_{24} through ω_{31} alone it is the case that $\omega(p, q) = \omega(q, pq) = \omega(pq, p)$ in the cases where

1. $p = 2r > 0, q = 1$
2. $p = 2r + 1, q = 1$
3. $p = 2r > 0, q = 2r + 1$
4. $p = 2r + 1, q = 2r + 1$

8.2. **The inductive step.** So now let us suppose that for $0 \neq r \neq s \neq 0$ it is true that $\omega(r, s) = \omega(s, rs) = \omega(rs, r)$ and try to establish the four inductive steps

1. $p = 2r, q = 2s$

   We will attempt to show that $\omega(2r, 2s) = \omega(2s, 2rs) = \omega(2rs, 2r)$. $\omega(2r, 2s) = -\omega(r, s)$ for ω_0 through ω_3 and ω_{24} through ω_{27}.

   $\omega(2r, 2s) = \omega(r, s)$ for ω_4 through ω_7 and ω_{28} through ω_{31}.

   $\omega(2s, 2rs) = -\omega(s, rs)$ for ω_0 through ω_3 and ω_{24} through ω_{27}. $\omega(2s, 2rs) = \omega(s, rs)$ for ω_4 through ω_7 and ω_{28} through ω_{31}.

$\omega(2rs, 2r) = \omega(rs, r)$ for ω_0 through ω_3 and ω_{24} through ω_{27}.
$\omega(2rs, 2r) = \omega(rs, r)$ for ω_4 through ω_7 and ω_{28} through ω_{31}.

Thus $\omega(2r, 2s) = \omega(2s, 2rs) = \omega(2rs, 2r)$ for ω_0 through ω_7 and ω_{24} through ω_{31}.

(2) $p = 2r, q = 2s + 1$
We will attempt to show that $\omega(2r, 2s+1) = \omega(2s+1, 2rs+1) = \omega(2rs + 1, 2r)$.
$\omega(2r, 2s + 1) = \omega(r, s)$ for $\omega_0, \omega_1, \omega_4, \omega_5, \omega_{24}, \omega_{25}, \omega_{28}, \omega_{29}$.
$\omega(2r, 2s + 1) = -\omega(r, s)$ for $\omega_2, \omega_3, \omega_6, \omega_7, \omega_{26}, \omega_{27}, \omega_{30}, \omega_{31}$.

$\omega(2s+1, 2rs+1) = \omega(s, rs)$ for $\omega_0, \omega_2, \omega_4, \omega_6, \omega_{24}, \omega_{26}, \omega_{28}, \omega_{30}$.
$\omega(2r, 2s + 1) = -\omega(s, rs)$ for $\omega_1, \omega_3, \omega_5, \omega_7, \omega_{25}, \omega_{27}, \omega_{29}, \omega_{31}$.

So this property fails for $\omega_1, \omega_2, \omega_5, \omega_6, \omega_{25}, \omega_{26}, \omega_{29}, \omega_{30}$.
$\omega(2rs + 1, 2r) = \omega(r, s)$ for $\omega_0, \omega_4, \omega_{24}, \omega_{28}$.
$\omega(2rs + 1, 2r) = -\omega(r, s)$ for $\omega_3, \omega_7, \omega_{27}, \omega_{31}$.
Thus $\omega(2r, 2s + 1) = \omega(2s + 1, 2rs + 1) = \omega(2rs + 1, 2r)$ for $\omega_0, \omega_3, \omega_4, \omega_7, \omega_{24}, \omega_{27}, \omega_{28}, \omega_{31}$.

(3) $p = 2r + 1, q = 2s$
The result $\omega(2r + 1, 2s) = \omega(2s + 1, 2rs + 1) = \omega(2rs + 1, 2r)$ follows immediately from $\omega(2r, 2s + 1) = \omega(2s + 1, 2rs + 1) = \omega(2rs + 1, 2r)$ and the fact that $\omega(p, q) = -\omega(q, p)$ for $0 \neq p \neq q \neq 0$.

(4) $p = 2r + 1, q = 2r + 1$
We will attempt to show that $\omega(2r+1, 2s+1) = \omega(2s+1, 2rs) = \omega(2rs, 2r + 1)$ for $\omega_0, \omega_3, \omega_4, \omega_7, \omega_{24}, \omega_{27}, \omega_{28}, \omega_{31}$.
$\omega(2r + 1, 2s + 1) = \omega(r, s)$ for $\omega_0, \omega_4, \omega_{24}, \omega_{28}$.
$\omega(2r + 1, 2s + 1) = -\omega(r, s)$ for $\omega_3, \omega_7, \omega_{27}, \omega_{31}$.

$\omega(2s + 1, 2rs) = \omega(s, rs)$ for $\omega_0, \omega_4, \omega_{24}, \omega_{28}$.
$\omega(2s + 1, 2rs) = \omega(s, rs)$ for $\omega_3, \omega_7, \omega_{27}, \omega_{31}$.

$\omega(2rs, 2r + 1) = \omega(rs, r)$ for $\omega_0, \omega_4, \omega_{24}, \omega_{28}$.
$\omega(2rs, 2r + 1) = -\omega(rs, r)$ for $\omega_3, \omega_7, \omega_{27}, \omega_{31}$.

Thus the quaternion properties are satisfied only by the eight twists $\omega_0, \omega_3, \omega_4, \omega_7, \omega_{24}, \omega_{27}, \omega_{28}, \omega_{31}$. Their twist constants are given in Table 5 on the next page. We notice that what distinguishes these eight from the remaining 24 is that $\alpha' = -\alpha$ and that $\gamma' = \gamma$. Only in such cases will the product satisfy the quaternion properties.

Table 7 on the facing page shows all eight twist tables for the octonions. To obtain the traditional quaternion sub-table, one must let $\alpha = 1$.

	α'	α	β	γ	γ'
$P_0 : (a,b)(c,d) = (ca - b^*d, da^* + bc)$	$-$	$+$	$-$	$+$	$+$
$P_3 : (a,b)(c,d) = (ca - db^*, a^*d + cb)$	$-$	$+$	$-$	$-$	$-$
$P_4 : (a,b)(c,d) = (ac - b^*d, da^* + bc)$	$-$	$+$	$+$	$+$	$+$
$P_7 : (a,b)(c,d) = (ac - db^*, a^*d + cb)$	$-$	$+$	$+$	$-$	$-$
$P_{24} : (a,b)(c,d) = (ca - bd^*, ad + c^*b)$	$+$	$-$	$-$	$+$	$+$
$P_{27} : (a,b)(c,d) = (ca - d^*b, da + bc^*)$	$+$	$-$	$-$	$-$	$-$
$P_{28} : (a,b)(c,d) = (ac - bd^*, ad + c^*b)$	$+$	$-$	$+$	$+$	$+$
$P_{31} : (a,b)(c,d) = (ac - d^*b, da + bc^*)$	$+$	$-$	$+$	$-$	$-$

Table 5. Twist Constants for the Cayley-Dickson Products



$\mathbf{C}$
$\mathbf{C}$ $\mathbf{T}$ $\mathbf{L}$ $-\mathbf{D}$
$\mathbf{L}$
$\mathbf{L}$ $\alpha\mathbf{N}$ $\mathbf{L}$ $-\alpha\mathbf{N}$
$\mathbf{T}$
$\mathbf{T}$ $\mathbf{T}$ $-\alpha\mathbf{N}$ $\alpha\mathbf{N}$
$-\mathbf{D}$
$-\mathbf{D}$ $-\alpha\mathbf{N}$ $\alpha\mathbf{N}$ $-\mathbf{D}$
$\mathbf{N}$
$\beta\mathbf{N}$ $\gamma\mathbf{N}$ $\gamma\mathbf{N}$ $\gamma\mathbf{N}$
$-\mathbf{N}$
$-\beta\mathbf{N}$ $-\gamma\mathbf{N}$ $-\gamma\mathbf{N}$ $-\gamma\mathbf{N}$

Table 6. Generalized Cayley-Dickson (actual) Twist Tree

9. The product P_0 and Octonion Index Cycling

In Cayley-Dickson spaces, if $i_p i_q = i_r$ then that fact is abbreviated (p, q, r). This is called a structure constant.

If $0 \neq p \neq q \neq 0$ then for all 32 products one and only one of (p, q, pq) or (q, p, pq) is true. For those eight which satisfy the quaternion property, (p, q, pq) implies (q, pq, p) which implies $(pq, p, q.)$

	0	1	2	3	4	5	6	7
0	1	1	1	1	1	1	1	1
1	1	-1	α	$-\alpha$	α	$-\alpha$	α	$-\alpha$
2	1	$-\alpha$	-1	α	$\alpha\beta$	$\alpha\gamma$	$-\alpha\beta$	$-\alpha\gamma$
3	1	α	$-\alpha$	-1	$\alpha\gamma$	$\alpha\gamma$	$-\alpha\gamma$	$-\alpha\gamma$
4	1	$-\alpha$	$-\alpha\beta$	$-\alpha\gamma$	-1	α	$\alpha\beta$	$\alpha\gamma$
5	1	α	$-\alpha\gamma$	$-\alpha\gamma$	$-\alpha$	-1	$\alpha\gamma$	$\alpha\gamma$
6	1	$-\alpha$	$\alpha\beta$	$\alpha\gamma$	$-\alpha\beta$	$-\alpha\gamma$	-1	α
7	1	α	$\alpha\gamma$	$\alpha\gamma$	$-\alpha\gamma$	$-\alpha\gamma$	$-\alpha$	-1

TABLE 7. Octonion Twist Table for the eight Cayley-Dickson products

	0	1	2	3	4	5	6	7
0	1	1	1	1	1	1	1	1
1	1	-1	1	-1	1	-1	1	-1
2	1	-1	-1	1	β	γ	$-\beta$	$-\gamma$
3	1	1	-1	-1	γ	γ	$-\gamma$	$-\gamma$
4	1	-1	$-\beta$	$-\gamma$	-1	1	β	γ
5	1	1	$-\gamma$	$-\gamma$	-1	-1	γ	γ
6	1	-1	β	γ	$-\beta$	$-\gamma$	-1	1
7	1	1	γ	γ	$-\gamma$	$-\gamma$	-1	-1

TABLE 8. Octonion Twist Table when $\alpha = 1$

C	**L**	**T**	**D**	**N**
$r = s = 0$	$r > s = 0$	$s > r = 0$	$r = s \neq 0$	$0 \neq r \neq s \neq 0$
$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} 1 & 1 \\ 1 & -1 \end{bmatrix}$	$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix}$	$\begin{bmatrix} \beta & \gamma \\ \gamma & \gamma \end{bmatrix}$

TABLE 9. Twist Table Components for $\omega_0, \omega_3, \omega_4, \omega_7$

If p is an integer and $1 \leq p \leq 7$ and π_0 is the permutation (1376524), then define the π_0 *successor* p' of p as the integer following p in the permutation. Then for the product P_0 alone it is true that (p, q, r) implies (p', q', r').

$$\begin{aligned}(1,2,3) &\to (3,4,7)\\ &\to (7,1,6)\\ &\to (6,3,5)\\ &\to (5,7,2)\\ &\to (2,6,4)\\ &\to (4,5,1)\\ &\to (1,2,3)\end{aligned}$$

This, together with the second quaternion property is sufficient to recover the P_0 multiplication table for the octonions.

1	i_1	i_2	i_3	i_4	i_5	i_6	i_7
i_1	-1	i_3	$-i_2$	i_5	$-i_4$	i_7	$-i_6$
i_2	$-i_3$	-1	i_1	$-i_6$	i_7	i_4	$-i_5$
i_3	i_2	$-i_1$	-1	i_7	i_6	$-i_5$	i_4
i_4	$-i_5$	i_6	$-i_7$	-1	i_1	$-i_2$	i_3
i_5	i_4	$-i_7$	$-i_6$	$-i_1$	-1	i_3	i_2
i_6	$-i_7$	$-i_4$	i_5	i_2	$-i_3$	-1	i_1
i_7	i_6	i_5	i_4	$-i_3$	$-i_2$	$-i_1$	-1

TABLE 10. Basis Vector Multiplication Table for P_0

10. Recursive definition of structure constants for the Cayley-Dickson twists

For $\omega_0, \omega_3, \omega_4$ and ω_7,

$$(1, 2n, 2n+1) \text{ for all } n > 0 \tag{35}$$

whereas for $\omega_{24}, \omega_{27}, \omega_{28}$ and ω_{31},

$$(2n, 1, 2n+1) \text{ for all } n > 0 \tag{36}$$

And for all eight of the Cayley-Dickson twists, the second quaternion property holds.

For $0 \neq p \neq q \neq 0$

$$(p, q, r) \longrightarrow (q, r, p) \longrightarrow (r, p, q) \tag{37}$$

For ω_0 and ω_{24} if $0 \neq p \neq q \neq 0$ then

$$\begin{aligned}(p,q,r) &\longrightarrow (2q,2p,2r)\\ &\longrightarrow (2p,2q+1,2r+1)\\ &\longrightarrow (2p+1,2q,2r+1)\\ &\longrightarrow (2p+1,2q+1,2r)\end{aligned}\tag{38}$$

For ω_3 and ω_{27} if $0 \neq p \neq q \neq 0$ then

$$\begin{aligned}(p,q,r) &\longrightarrow (2q,2p,2r)\\ &\longrightarrow (2q,2p+1,2r+1)\\ &\longrightarrow (2q+1,2p,2r+1)\\ &\longrightarrow (2q+1,2p+1,2r)\end{aligned}\tag{39}$$

For ω_4 and ω_{25} if $0 \neq p \neq q \neq 0$ then

$$\begin{aligned}(p,q,r) &\longrightarrow (2p,2q,2r)\\ &\longrightarrow (2p,2q+1,2r+1)\\ &\longrightarrow (2+1,2r,2r+1)\\ &\longrightarrow (2p+1,2q+1,2r)\end{aligned}\tag{40}$$

For ω_7 and ω_{31} if $0 \neq p \neq q \neq 0$ then

$$\begin{aligned}(p,q,r) &\longrightarrow (2p,2q,2r)\\ &\longrightarrow (2q,2p+1,2r+1)\\ &\longrightarrow (2q+1,2p,2r+1)\\ &\longrightarrow (2q+1,2p+1,2r)\end{aligned}\tag{41}$$

11. Octonion Cycles

If $0 \neq p \neq q \neq 0$, and if $i_p i_q = +i_r$ we say that the *sense* of (p,q,r) (and therefore of (q,r,p) and $r,p,q)$) is *positive* or that $\sigma(p,q,r) = 1$. Otherwise we say that the the sense is *negative* and $\sigma(p,q,r) = -1$.

So it follows that for the twist constants α, β and γ

$$\alpha = \sigma(1,2,3)\tag{42}$$

$$\alpha\beta = \sigma(2,4,6)\tag{43}$$

$$\alpha\gamma = \sigma(2,5,7)\tag{44}$$

If we define π as the permutation $(1)(357)(246)$ and for $1 \leq p \leq 7$ we let p'' denote the π successor of p, then

$$\sigma(p,q,r) = \sigma(p'',q'',r'') \tag{45}$$

Thus, for any of the eight Cayley-Dickson doubling variations, knowing the sense of $(1,2,3)$, $(2,4,6)$ and $(2,5,7)$ is sufficient to recover the sense of all 42 triples (p,q,r).

One way to recover any of the eight variations on the sets of 42 triples is by using the Fano plane (Figure 1 on the next page).

Each of the three sides of the triangle represents a triple (p,q,r). So does each altitude from a vertex to the midpoint of the opposite side and so does the circle through the midpoints. For each of the eight Cayley-Dickson products, the sense of the three sides is the same–either clockwise ($\leftarrow$) around the triangle, or counter-clockwise ($\rightarrow$). If clockwise, then the three sides of the triangle represent $(5,2,7)$, $(7,4,3)$ and $(3,6,5)$. If counter-clockwise then the three sides represent $(7,2,5)$, $(5,6,3)$ and $(3,4,7)$. The circle through the midpoints of the sides represents either $(2,4,6)$ in the clockwise sense ($\circlearrowright$) or $(2,6,4)$ in the counter-clockwise sense ($\circlearrowleft$). The three altitudes may all be in an 'up' sense from center to vertex ($\uparrow$) or may all be in a 'down' sense from center to base ($\downarrow$). So the altitudes must either be $(2,1,3)$, $(6,1,7)$, and $(4,1,5)$ or $(3,1,2)$, $(7,1,6)$, and $(5,1,4)$. All altitudes must have the same sense. See Table 11 on the facing page for a breakdown of all the modes of the eight Cayley-Dickson doubling products. A thing to notice about the Fano plane is that a 120° counter-clockwise rotation of the diagram represents the permutation $\pi = (1)(357)(246)$ mentioned above.

Since the sides may have two senses and the circle may have two senses and the altitudes may have two senses, all $2^3 = 8$ versions of the Cayley-Dickson products may be accommodated in the one diagram.



Figure 1. Fano Plane

$P_0 : (a,b)(c,d) = (ca - b^*d, da^* + bc)$	$\downarrow$	$\circlearrowleft$	$\rightarrow$
$P_3 : (a,b)(c,d) = (ca - db^*, a^*d + cb)$	$\downarrow$	$\circlearrowleft$	$\leftarrow$
$P_4 : (a,b)(c,d) = (ac - b^*d, da^* + bc)$	$\downarrow$	$\circlearrowright$	$\rightarrow$
$P_7 : (a,b)(c,d) = (ac - db^*, a^*d + cb)$	$\downarrow$	$\circlearrowright$	$\leftarrow$
$P_{24} : (a,b)(c,d) = (ca - bd^*, ad + c^*b)$	$\uparrow$	$\circlearrowright$	$\leftarrow$
$P_{27} : (a,b)(c,d) = (ca - d^*b, da + bc^*)$	$\uparrow$	$\circlearrowright$	$\rightarrow$
$P_{28} : (a,b)(c,d) = (ac - bd^*, ad + c^*b)$	$\uparrow$	$\circlearrowleft$	$\leftarrow$
$P_{31} : (a,b)(c,d) = (ac - d^*b, da + bc^*)$	$\uparrow$	$\circlearrowleft$	$\rightarrow$

Table 11. Fano Plane Modes

References

[1] W. Ambrose *Structure theorems for a special class of Banach algebras* Trans. Amer. Math. Soc. Vol. **57**, (1945) 364-386

[2] J. Baez *The Octonions* Bul. Am. Math. Soc. Vol. **39**, No. 2, (2001) 145-205

[3] J. Bales *Properly twisted groups and their algebras* (2006) arXiv:1107.1297

[4] J. Bales *A tree for computing the Cayley-Dickson twist* Missouri J. of Math. Sci. Vol 21 No. 2 (2009)

[5] M. Bremner, I. Hentzel *Identities for algebras obtained from the Cayley-Dickson process* Comm. Algebra Vol. **29** (2001) 3523-3534

[6] R. Brown, *On generalized Cayley-Dickson algebras* Pacific J. Math. Vol. **20** (1967)

[7] R. Busby, H. Smith *Representations of twisted group algebras* Trans. Am. Math. Soc. Vol. **149**, No. 2, (1970) 503-537

[8] G. Dixon *Octonion X-product Orbits* arXiv:hep-th/9410202v1 26 Oct 1994

[9] R. Erdmann *Über verallgemeinerte Cayley-Dickson Algebren* J. Reine Angew. Math. Vol. **250** (1971) 153-181

[10] N. Jacobson, *Basic Algebra I*, W. H. Freeman, San Francisco, 1974, 417-427

[11] K. McCrimmon *Derivations and Cayley derivations of generalized Cayley-Dickson algebras* Pacific J. Math. Vol. **117** (1985) 163-182

[12] G. Moreno *The zero divisors of the Cayley-Dickson algebras over the reals* Bol. Soc. Mat. Mex. **4**, (1998) 13-27

[13] R. Schafer *On the algebras formed by the Cayley-Dickson process* Amer. J. Math. Vol. **76** (1954) 435-446

DEPARTMENT OF MATHEMATICS, TUSKEGEE UNIVERSITY, TUSKEGEE, AL 36088, (JUNE 2011)

E-mail address: `jbales@mytu.tuskegee.edu`